

**The Dichotomus Horizon:  
proposal for a new biochronologic unit of the Giraudi Zone  
of the Upper Barremian of southeastern France,  
and considerations regarding the genus *Imerites* ROUCHADZÉ  
(Ammonoidea, Gassendiceratinae)**

Didier BERT <sup>1</sup>

G rard DELANOY <sup>2</sup>

St phane BERSAC <sup>3</sup>

**Abstract:** Recent revisions of the genus *Imerites* ROUCHADZ  make it possible to introduce a new biochronologic horizon to define more precisely the lower boundary of the Giraudi Zone: the Dichotomus Horizon. Using the concept of 'interval zone', this new horizon maintains the current lower boundary of the Giraudi Zone as accepted by authors, and thus contributes to the stabilization of the Barremian zonal system. This stabilization is also strengthened by abandonment of the use of "*Crioceras*" *cristatus* d'ORBIGNY (*nomen dubium*) that ought not be used as an index species in detriment of *Imerites giraudi* (KILIAN). The classification, origin, and intraspecific variation of the genus *Imerites* ROUCHADZ  are examined.

**Key Words:** Tethyan realm; Upper Barremian; biostratigraphy; Interval zone; southeastern France; ammonites.

**Citation:** BERT D., DELANOY G. & BERSAC D. (2011).- The Dichotomus Horizon: proposal for a new biochronologic unit of the Giraudi Zone of the Upper Barremian of southeastern France, and considerations regarding the genus *Imerites* ROUCHADZ  (Ammonoidea, Gassendiceratinae).- [Carnets de G ologie / Notebooks on Geology](#), Brest, Article 2011/01 (CG2011\_A01)

**R sum  :** *L'horizon   Dichotomus : proposition d'une nouvelle unit  biochronologique de la zone   Giraudi du Barr mien sup rieur du Sud-Est de la France, et consid rations sur le genre Imerites ROUCHADZ  (Ammonitina, Gassendiceratinae).*- Les r centes r visions du genre *Imerites* ROUCHADZ  permettent   pr sent d'introduire un nouvel horizon biochronologique afin de mieux d finir la limite inf rieure de la Zone   Giraudi : l'horizon   Dichotomus. Par l'application du concept de la zone d'intervalle, l'utilisation de cet horizon permet de conserver la limite inf rieure actuelle de la zone   Giraudi telle qu'elle a  t  accept e par les auteurs, et ainsi de contribuer   la stabilisation du sch ma zonal du Barr mien. Cette stabilisation est aussi renforc e par l'abandon de l'usage de "*Crioceras*" *cristatus* d'ORBIGNY (*nomen dubium*) qui ne devrait pas  tre utilis  comme esp ce indice au d triment de *Imerites giraudi* (KILIAN). La classification, l'origine et la variabilit  intrasp cifique du genre *Imerites* ROUCHADZ  sont aussi discut es.

**Mots-Clefs :** Domaine t thysien ; Barr mien sup rieur ; biostratigraphie ; zone d'intervalle ; Sud-Est de la France ; ammonites.

## Introduction

After a period of instability, in recent years several studies have helped to fix the calibration of the Upper Barremian biostratigraphy of southeastern France that increased its value for practical use and increased the degree of its re-

producibility by specialists working in discrete localities (REBOULET *et alii*, 2007, 2009 - see BERT *et alii*, 2008 for a historical account). It now includes the Vandenheckei, Sartousiana and Giraudi zones (Fig. 1), their limits based on faunal changes commonly associated with sequence boundaries developed in relation to

---

<sup>1</sup> \* Corresponding author

Universit  de Bourgogne, Laboratoire Biog osciences, UMR CNRS 5561, 6 bd Gabriel, F-21000, Dijon (France)  
[paleo-db@orange.fr](mailto:paleo-db@orange.fr)

<sup>2</sup> D partement des Sciences de la Terre, Universit  de Nice-Sophia-Antipolis, Facult  des Sciences, 28 avenue Valrose, 06108 Nice Cedex 2 (France)  
[delanoy@unice.fr](mailto:delanoy@unice.fr)

<sup>3</sup> 945 route de Gatti res, F-06640 Saint Jeannet (France)  
[geosteber@yahoo.fr](mailto:geosteber@yahoo.fr)

Manuscript online since January 17, 2011

eustasy (ARNAUD, 2005). In accord with the recommendations of the *IUGS Lower Cretaceous Ammonite Working Group, the KILIAN Group*, stabilization of the boundaries of these zones led to a preference for the use of interval zones rather than zones based on the range of a taxon. With good reason, for the boundaries of zones based on individual ranges are subject to repeated change in range limits caused by new finds or by tergiversation resulting from a revision of index taxa. Thus it seems preferable to choose stable and precise horizons for defining the base of zones (THIERRY, 1997). The introduction of several biochronological horizons

(DELANOY, 1995, 1997, 1998; BERT *et alii*, 2008; BERT & DELANOY, 2009; BERT *et alii*, 2010) has contributed significantly to a refinement of the biostratigraphic pattern of the Tethyan Upper Barremian of southeastern France.

This work revises the definition of the lower boundary of the Giraudi Zone [index species: *Imerites giraudi* (KILIAN, 1888)] and is a manifestation of progress in the continuation of the revision of Upper Barremian biostratigraphy in southeastern France through emendation to the ammonite faunas which are a major element of its framework (here *Imerites* ROUCHADZÉ, 1933).

Stages	Reboulet <i>et alii</i> , 2009			This work			
	Zones	Sub-Zones	Horizons	Zones	Sub-Zones	Horizons	
Upper Barremian	Giraudi	Waagenoides		Giraudi	Waagenoides		
		Sarasini			Sarasini		
			Puzosianum			Puzosianum	
		Giraudi	Emerici		Giraudi	Giraudi	Emerici
			Giraudi				Giraudi
			Dichotomus				Dichotomus
	Sartousiana	Feraudianus		Sartousiana	Feraudianus	Autrani	
		Provincialis			Provincialis	Bersaci	
						Magnini	
						Feraudianus	
		Sartousiana			Limentinus	Casanovai	
		Vandenheckei	Barremense			Vandenheckei	Alpinum
				Sartousiana			
				Limentinus			
	Vandenheckei			Vandenheckei	Vandenheckei		Marchandi
			Breistrofferi				
			Alpinum				


◀ **Figure 1:** Zonal scheme amended after BERT *et alii*, 2008, and REBOULET *et alii*, 2009. In red the Dichotomus Horizon (new). **Figure 1 :** Schéma zonal modifié d'après BERT *et alii*, 2008, et REBOULET *et alii*, 2009. En rouge, l'horizon à *Dichotomus* (nouveau).

### The boundary between the Sartousiana and Giraudi zones, and the new Dichotomus Horizon

#### 1- Geological setting

The Lower Cretaceous of southeastern France is marked by the evolution of a large intracratonic subsident area known as the Vocontian Basin (PAQUIER, 1900). The area of the historical Barremian stratotype (Angles-Barrême-Castellane area – Fig. 2), is in the southern part of the Vocontian Basin which is less affected by gravity remodelling and Alpine orogeny than its

northern portion. The Barremian here is characterized by pelagic sediments, mainly alternation of marls and limestones in decimetric to metric beds. Given the relative continuity of deposits and the paleontological record, it is possible to track in considerable detail the succession and evolution of their ammonite faunas. Thus, following up on the work of DELANOY (1995, 1998), several sections of this area expose the boundary between the Sartousiana and the Giraudi zones well enough that they can be examined minutely: they are the Vignon section (VIG, Fig. 3), the Descouère section (DES, Fig. 4), and the Grande-Terre section (GT, Fig. 5).



◀ **Figure 2:** Barremian paleogeography of southeastern France and location of the area studied (from ARNAUD, 2005, amended). **Figure 2 :** Cadre paléogéographique du Sud-Est de la France et localisation du secteur d'étude (modifiée d'après ARNAUD, 2005).

## 2- The boundary between the Sartousiana and Giraudi zones in southeastern France

The base of the Giraudi Zone is characterized by an important phase of marine transgression (ARNAUD, 2005), marked lithologically by the "vire marneuse à *Heteroceras*" of authors. It is associated with a major faunal turnover (DELANOY, 1990, 1995, 1997, 1998; BERT *et alii*, 2008). This turnover is relatively progressive in strata at the top of the Feraudianus Subzone and at the base of the Giraudi Subzone. In fact, all change occurs between the lower part of the Feraudianus Subzone and the Emerici Horizon of the Giraudi Subzone (Fig. 1) where the Hemihoplitinae are progressively replaced quantitatively by the Heteroceratidae. At the base of the Feraudianus Subzone Hemihoplitinae are the major components of the ammonite fauna with a preponderance of the genus *Hemihoplites* SPATH. The Pulchelliidae and the Peircinae are quite rare, as are the Gassendiceratinae (genera *Gassendiceras* BERT, DELANOY & BERSAC, 2006, and *Pseudoshasticioceras* DELANOY, 1998) which become more numerous at the top of the Subzone (*i.e.* in the Bersaci and Autrani horizons). The genus *Heteroceras* d'ORBIGNY (with a turruculate morphology) is present at the upper limit of the Feraudianus

Subzone (Autrani Horizon) but is extremely rare, for at this level and time it is only a very minor element of the Giraudi fauna. In the lower portion of the Giraudi Subzone the situation is reversed: the Hemihoplitinae and the Pulchelliidae are gone and turruculate morphology dominates quantitatively, first briefly with *Imerites* (Gassendiceratinae), and later with *Heteroceras* which begin an increase in abundance at the base of the Giraudi Zone and proliferate accompanied by a morphological explosion in the Emerici Horizon (DELANOY, 1990, 1995, 1997, 1998; DELANOY & EBBO, 2000; DELANOY & BERT, 2006).

Above the Autrani Horizon (Feraudianus Subzone, Sartousiana Zone, Fig. 1), the appearance of the genus *Imerites* ROUCHADZÉ is currently accepted by authors as the valid marker of the lower limit of the Giraudi Zone (see historical account in KAKABADZE, 1989; HOEDEMAEKER & BULOT, 1990; DELANOY, 1990, 1995, 1998; REBOULET *et alii*, 2006, 2007, 2009; BERT *et alii*, 2008). Revisions of this genus by DELANOY (1998) and BERT *et alii* (2009), improved understanding of the stratigraphic distribution of the species of *Imerites*. Their development over time is coincident with the evolutionary framework of the last Gassendiceratinae BERT, DELANOY & BERSAC, 2006. So in the succession

*Imerites dichotomus* ERISTAVI, 1955 appears before *Imerites giraudi* (KILIAN, 1888). The use of an interval zone allows a redefinition of the lower boundary of the Giraudi Zone without change of level by use of the new Dichotomus Horizon (defined below). This usage preserves the integrity and stability of Barremian zonation (BERT *et alii*, 2008). There is no need to move the lower limit of the Giraudi Zone upward to make it coincident with the appearance of its index species, or to rename the Giraudi Zone as the Dichotomus Zone, because *I. dichotomus* ERISTAVI is the first species of *Imerites* of which the appearance coincides with the accepted definition of the lower limit of the Giraudi Zone. In any event, any shift in level would dissociate the base of the Giraudi Zone from the faunal turnover and the sequence boundary with which it is coincident (ARNAUD, 2005); application either of these options would change the level of the base of the zone and return to the instability that has long characterized Barremian zonation and should cease.

### 3- The Dichotomus Horizon (*new*)

In France *Imerites dichotomus* ERISTAVI has a very limited and precise stratigraphic position. By anagenesis (BERT *et alii*, 2009) it is the successor of the index species *Pseudoshasticrioceras autrani* (DELANOY), and the ancestor of the index species *Imerites giraudi* (KILIAN). In the stratotype area (Vocontian Basin, southeastern France – Fig. 2) its appearance in the stratigraphic succession is in agreement with its biologic relationships. These facts, and the need for the establishment of a high resolution biostratigraphy for the whole of the Barremian (REBOULET *et alii*, 2006; REBOULET *et alii*, 2007; BERT *et alii*, 2008; REBOULET *et alii*, 2009), has impelled us to propose *Imerites dichotomus* ERISTAVI as a new biostratigraphic marker in the Vocontian Basin. This species occurs in strata immediately above the major beds of the Autrani Horizon at the top of the Feraudianus Subzone (Sartousiana Zone), and immediately precedes those of the Giraudi Horizon where *Imerites giraudi* (KILIAN) occurs (Figs. 3 - 4 - 5).

**Index species:** *Imerites dichotomus* ERISTAVI, recently revised by BERT *et alii* (2009).

**Status:** This horizon is defined by the first appearance of its index species (bed No. 436 in the Vignon [VIG] section, Fig. 3), and its upper limit is currently set at the base of the Giraudi Horizon (bed No. 439 in the VIG section) with the first appearance of *Imerites giraudi* (KILIAN). The Dichotomus Horizon is also present in the sections near La Baume (Castellane area): beds 151 to 152 in the Descouère section (DES, Fig. 4), and bed 679 in the Grande-Terre section (GT, Fig. 5) [see also DELANOY, 1995, 1998].

**Paleobiogeographic distribution:** *Imerites dichotomus* ERISTAVI is present in south-eastern France but also in Spain (oral commu-

nication of COMPANY, see DELANOY, 1998, p. 207), Bulgaria, Romania and Georgia (see BERT *et alii*, 2009), a distribution that augers an extensive application of the Dichotomus Horizon.

**Faunal assemblages:** The index species is generally fairly well represented in the French sections. It is associated with (see DELANOY, 1995, 1998; BERT *et alii*, 2008): *Macroscaphites yvani* (PUZOS) macro- and microconchs, *Acantholytoceras pseudoaudouli* (THOMEL) macro- and microconchs, *Jaubertites collignoni* SARKAR, *Protetragonites crebrisulcatus* (UHLIG), *Eulytoceras phestus* (MATHERON), *Silesites seranonis* (d'ORBIGNY), *Melchiorites melchioris* (TIETZE), *Barremites difficilis* (d'ORBIGNY), *Barremites strettostoma* (UHLIG), *Phylloceras ponticuli* (ROUSSEAU), and with *Heteroceras couletti* DELANOY, *Heteroceras baylei* REYNES and *Spinocrioceras trachyomphalus* (UHLIG).


## Considerations regarding the genus *Imerites* ROUCHADZÉ, 1933

### 1- Remarks on the classification and origin of *Imerites*

The genus *Imerites* ROUCHADZÉ has traditionally been classified as a Heteroceratidae, but it is now recognized as a representative of the family Hemihoplitidae (see BERT *et alii*, 2009 for an historical account) despite the presence of a turricon in the juvenile part of the shell (helicoidal coiling). SARKAR was the first to classify *Imerites* (his *Escragnoileites*) in the Hemihoplitidae (1955, p. 24) because it has a hemihoplitid ancestor (1955, p. 18, 22), although KILIAN (1907-1913) saw "some similarities" between *Heteroceras giraudi* (= *Imerites*) and some Hemihoplitidae [= *Ancyloceras* (*Crioceras*) *heberti* in KILIAN's time]. It has been recognized for a long time that helicoidal coiling is not restricted to the Heteroceratidae. Indeed, KILIAN (1888, 1889) was the first to restrict the genus *Heteroceras* d'ORBIGNY only to those Barremian taxa that have a turricon. But d'ORBIGNY classified one Senonian species (1851, p. 222) as a heteroceratid, and MEEK (1876) included helicoidal forms from the Upper Cretaceous of the United States of America (Nostoceratidae) in that group, thus indicating broader criteria for their classifications. This type of coiling is known to have developed repeatedly in the evolutionary history of the ammonoids (e.g. Triassic *Cochloceras*; some Bajocian *Spiroceras*; Uppermost Barremian *Kutatissites*; some Lower Cretaceous Leptoceratoidae; Albian *Mariella*, *Turrillitoides*, *Helicoceras* or *Pseudhelicoceras*; Cenomanian *Turrillites*, *Hypoturrillites*, *Mesoturrillites* and *Ostlingoceras*; Axonoceras, *Jouaniceras*, *Anaklinoceras* and some other Nostoceratidae from the Upper Cretaceous, etc. – see ARKELL *et alii*, 1957 and WRIGHT *et alii*, 1996). But these authors did not remark a connection between these homeomorphs and the Heteroceratidae. More recently, KAKABADZE (2004, p.

21) made the following observation about certain genera with helicoidal coiling and the absence of any relationship to the Heteroceratidae: "the similarity in the mode of coiling (helicoidal, planispiral or helicoidal, planispiral and uncoiled) is not important for identifying

the systematic position as to family". And DELANOY (1998, p. 184) said about *Imerites* that "the presence of the turricone would be only the expression of a homeomorphism that affects the early developments of these forms" (translation pars).



◀ **Figure 3:** Distribution of the ammonites in the Vignon (VIG) section *pars* (Barrême area, Alpes-de-Haute-Provence).

**Figure 3 :** Répartition des faunes d'ammonites dans la coupe du Vignon (VIG) *pars* (secteur de Barrême, Alpes-de-Haute-Provence).



BERT *et alii* (2009) were the first to distinguish and define evolutionary processes in the origin and development of species in the genus *Imerites* of the Hemihoplitidae. Their work involved study of an evolved "*Pseudoshastrioceras*" ornament in *Imerites dichotomus* ERISTAVI (see BERT *et alii*, 2006). In that stage of development weaker peri-ventral tubercles de-

monstrate that *Imerites* is a direct descendant of the genus *Pseudoshastrioceras* DELANOY (Gassendiceratinae BERT, DELANOY & BERSAC). This taxonomic distinction between genera of the same lineage is based on the early appearance of a turricone during the growth of *Imerites* (BERT *et alii* 2009).




Figure 4: Distribution of the ammonites in the Descouère (DES) section *pars* (La Baume in the Castellane area, Alpes-de-Haute-Provence).

Figure 4 : Répartition des faunes d'ammonites dans la coupe du Descouère (DES) *pars* (La Baume, secteur de Castellane, Alpes-de-Haute-Provence).





Figure 5: Distribution of the ammonites in the Grande-Terre (GT) section *pars* (La Baume in the Castellane area, Alpes-de-Haute-Provence).

Figure 5 : Répartition des faunes d'ammonites dans la coupe de la Grande-Terre (GT) *pars* (La Baume, secteur de Castellane, Alpes-de-Haute-Provence).

We welcome acceptance of the genus *Imerites* into the Gassendiceratinae but we do not agree with the broader generic limits proposed for *Imerites* by VERMEULEN & LEPINAY (2010, p. 18, 20). Their diagnosis includes some non-turriculate Hemihoplitidae, which would cause problems that preclude its acceptance:

1. the large range of variation in morphology and ornament proposed for their *Imerites* by these authors requires a redefinition of this genus to include criteria much broader in scope than those commonly acceptable to specialists. Their proposal widely depasses the intent of the author of the genus *Imerites* ROUCHADZÉ and that of subsequent workers. This new definition would produce a genus-group taxon which would greatly lessen the precision of identification and make correlation with previous work difficult. So nomenclatural stability would be threatened and Upper Barremian biostratigraphy again subject to revision because of confusion regarding the stratigraphic ranges of the species of the genus *Imerites* ROUCHADZÉ;
2. the ornamental and morphological characteristics specified in the emended diagnosis of VERMEULEN & LEPINAY, would include in their genus *Imerites* species of Hemihoplitinae currently assigned to other genera, although they lack any phyletic link to them. So *Imerites* in VERMEULEN & LEPINAY'S usage, might well include the taxa *Hemihoplites* SPATH, *Camereiceras* DELANOY, *Pachyhemihoplites* DELANOY or *Ancylezeiceras* VERMEULEN.

This problem is particularly evident in the taxon "*I.* *stephaniae* VERMEULEN & LEPINAY, 2010 (found in the Provincialis Subzone: One subzone separates it from the appearance of the first *Imerites* s. str.) that they attribute to the genus *Imerites* ROUCHADZÉ, but do not compare it with contemporary Hemihoplitids. "*I.* *stephaniae* VERMEULEN & LEPINAY is a fragmentary specimen (VERMEULEN & LEPINAY, 2010, p. 21, Pl. 1, fig. 7) particularly close in aspect to contemporary *Hemihoplites* of the group *H. casanovai* DELANOY (= *H. intermedius* VERMEULEN - currently under study).

## 2- The case of "*Crioceras*" *cristatus* d'ORBIGNY, 1842


AS DELANOY (in GAUTHIER *et alii*, 2006, p. 138-139), maintained "*Crioceras*" *cristatus* d'ORBIGNY should be considered synonymous with *Imerites giraudi* (KILIAN) and according to the principle of priority of the *International Code of Zoological Nomenclature* (ICZN - Art. 23.1) has seniority. Consequently, for VERMEULEN & LEPINAY (2010, p. 20) another amendment of Barremian zonation

would be necessary: renaming the Giraudi Zone as the *Cristatus* Zone [note that this is not obligatory, for the *International Stratigraphic Guide* (SALVADOR, 1994, p. 67) states that "if it is desirable to continue use of a taxonomic term which is no longer valid, the term should be in quotation marks"].

However new data obtained in the recent revision of *Imerites* by BERT *et alii* (2009, p. 32-33) found the syntypes of "*Crioceras*" *cristatus* d'ORBIGNY of the d'ORBIGNY'S collection are too fragmented to be identified at a specific level, and the fragments may not be of just one specimen but may even represent more than one species. These facts render this taxon unusable and a *nomen dubium*. Moreover, d'ORBIGNY'S (1842, Pl. 115, figs. 4-8) original illustrations are probably a synthetic compilations (*i.e.* a picture based on several different but fragmentary specimens assembled and restored to appear as if they were a single specimen). They were recognized as such by KILIAN (1888, 1889). In the original illustration the turricone is replaced by planispiral whorls for the stages of growth of the actual syntypes (particularly fragmented) when compared to the original picture make this obvious. So the rehabilitation of "*Crioceras*" *cristatus* d'ORBIGNY, 1842, to replace *Imerites giraudi* (KILIAN, 1888), a long-used, well defined taxon (recognized by many authors), is not desirable. Moreover, the purpose of the ICZN principle of priority is explained by Article 23.2 which states very explicitly that it must foster nomenclatural stability: "it is not intended to be used to upset a long-accepted name in its accustomed meaning by the introduction of a name that is its senior synonym or homonym, or through an action taken following the discovery of a prior and hitherto unrecognized nomenclatural act"; *a fortiori* when the older name is based on a synthetic compilations and can be considered as a *nomen dubium* because it differs from characteristics found in its syntypes. On the other hand, the proposal of invalidation by VERMEULEN & LEPINAY (2010, p. 17-18) of the d'ORBIGNY'S syntypes in favour to a more complete neotype (not among the syntypes) does not comply with the provisions of the ICZN (Art. 75, and in particular, § 75.3.4) and therefore cannot be implemented. At most it might be possible to designate one of the original syntypes as a lectotype (n°5405-1 of the d'ORBIGNY'S collection), but the basic problem remains unchanged...

Whatever the nomenclatural possibilities mentioned earlier (DELANOY in GAUTHIER *et alii*, 2006, p. 139), *Imerites giraudi* (KILIAN) is the type-species of the genus *Imerites* ROUCHADZÉ and we strongly recommend "*Crioceras*" *cristatus* d'ORBIGNY be considered invalid.





**Figure 6:** Dimorphism and variability in shell morphology in *Imerites*.  
**Figure 6 :** Dimorphisme et variabilité morphologique de la coquille chez *Imerites*.

### 3 - Intraspecific variability in *Imerites* and mechanical constraints associated with the presence of turricone

Intraspecific variability in species of *Imerites* involves ornamentation, some dimensional parameters of the shell and the size and orientation of the turricone (Fig. 6c). The relationship of turricone size and placement to changes in shell morphology has been studied by DELANOY (1998, p. 53) for the genus *Heteroceras* and was illustrated by KAKABADZE (2004, Fig. 10) for the *Kutatissites*. In *Imerites*, the phenomenon is the same: a larger turricone generally is accompanied by a less regular or triangular coiling before normal involution resumes. However, contrary to the views of VERMEULEN & LEPINAY (2010) these purely mechanical constraints are not comparable with the dimorphic differentiation in *Imerites dichotomus* ERISTAVI where tripartite adult microconch [m] are associated with large planospiral adult macroconch [M] (see BERT *et alii*, 2009, Pl. 2, fig. 2 *versus* Pl. 1, fig. 1 – Fig. 6a) for several reasons:

1. The dimorphs of the adult stages of *I. dichotomum* ERISTAVI are very different and there are, to our knowledge, no intermediaries (Fig. 6a);
2. The presence of such intermediaries was assumed by VERMEULEN & LEPINAY (2010) because of the existence of specimens with a triangular coil ("*I. cristatus favrei morphotype*" of Pl. 1, figs. 5-6). These specimens of which the whorl "may represent a draft shaft" (p. 18) are too small in size and incomplete. They are only juvenile whorls with triangular coiling that wraps the turricone, but lack an uncoiling of the outer whorls to form a shaft (Fig. 6c). Only the specimens figured Pl. 1, figs. 1-2 of their publication may be adult, but they are cryptoconic with no shaft. Consequently, no adult specimen with a morphology intermediate between those of micro- and macroconchs are known;
3. The tripartite *Imerites raricostatus* (KAKABADZE) has a well expressed ornamental stage with fibular ribs (the type specimen is refigured in KOTETISHVILI *et alii*, 2005, Pl. 87, fig. 2) and its ontogenetic sequence is very similar to the tripartite specimen figured in BERT *et alii* (2009, Pl. 2, fig. 2 – Fig. 6a). Therefore *I. raricostatus* (KAKABADZE) is a microconch of *I. dichotomus* ERISTAVI and not a morphotype of *I. giraudi* (KILIAN) (VERMEULEN & LEPINAY, 2010). Complete adult microconch specimens of *I. dichotomus*

ERISTAVI seem quite rare, but adult macroconchs are also few. The adult form of the macroconch was unknown prior to the recent revision by BERT *et alii* (2009). Juvenile or fragmentary specimens of this species that are difficult to place because of uncertainty regarding which dimorph they represent are relatively common;

4. Variability in the size of the turricone *versus* that of the morphology of the shell is well expressed in the macroconchs of *Imerites dichotomus* ERISTAVI, just as it is in *Imerites giraudi* (KILIAN) (Fig. 6c). This is the same in the microconchs of *I. dichotomus* (Fig. 6b): the type specimen of *I. rarecostatus* (= *I. dichotomus* [m]) has a large turricone and minute coil, whereas the specimen figured Pl. 2, fig. 2 (in BERT *et alii*, 2009) has a well-developed coil and a relatively small turricone;
5. The specimens of *I. giraudi* (= *I. cristatus* in VERMEULEN & LEPINAY, 2010) cannot be used to suggest the absence of dimorphism in *I. dichotomus*, since the tripartite coiling of the microconchs is currently recognized with certainty only in *I. dichotomus*.

### Conclusions

Recent revisions of the genus *Imerites* ROUCHADZÉ provide an increased understanding of its variability and its evolution, so now permit the introduction of a new biochronologic horizon, the *Dichotomus* Horizon (Fig. 1), to define the lower boundary of the Giraudi Zone more precisely. Using the interval zone concept, the addition of this horizon maintains the current lower boundary of the Giraudi Zone as accepted by authors (the appearance of the genus *Imerites*), and thus contributes to the stabilization of the Barremian zonal scheme. This stabilization is also strengthened by the abandonment of "*Crioceras*" *cristatus* d'ORBIGNY (*nomen dubium*) that ought not be used as index species instead of *Imerites giraudi* (KILIAN). Following THIERRY (1997), in general (as it is for the Barremian) it is preferable to use interval zones rather than the stratigraphic range of any one taxon for example, to avoid constant changes of boundaries that discoveries and systematic reviews of index taxa might impose, thus threatening the stability of the zonation. The introduction of a new biochronologic horizon increases the precision of the biostratigraphic zonation of the Barremian stage in southeastern France which is a requirement for the study in detail of the evolution of ammonites and their populations.

## Acknowledgments

We wish to express our thanks to Mr. Raymond ENAY, and to the anonymous reviewer, Mr. Stephane REBOULET and to Mr. Jaap KLEIN for their valuable advice. We thank the Réserve Géologique de Haute Provence and Mrs. Myette GUIOMAR who have allowed us access to the areas studied. One of us (D.B.) has interesting discussions with Gerd E.G. WESTERMANN about stratigraphical nomenclature, and we warmly thank him. We want especially and sincerely to thank Nestor SANDER who kindly corrected our English in the final version.

## Bibliographic references

- ARKELL W.J., KUMMEL B. & WRIGHT C.W. (1957).- Mesozoic Ammonoidea. *In*: MOORE R.C. (Ed.), Mollusca 4.- *Treatise on Invertebrate Paleontology*, Geological Society of America, New York; University of Kansas, Lawrence, Part L, p. L80-L437.
- ARNAUD H. (2005).- The South-East France Basin (SFB) and its Mesozoic evolution.- *Géologie Alpine*, Grenoble, (série spéciale "Colloques et Excursions"), n° 7, p. 5-28.
- BERT D. & DELANOY G. (2009).- *Pseudoshasticrioceras bersaci* nov. sp. (Ammonoidea, Gassendiceratinae), and new ammonite biohorizon for the Upper Barremian of south-eastern France.- *Carnets de Géologie / Notebooks on Geology*, Article 2009/02 (CG2009\_A02), 22 p.
- BERT D., DELANOY G. & BERSAC S. (2006).- Descriptions de représentants nouveaux ou peu connus de la Famille des Hemihoplitidae SPATH, 1924 (Barrémien supérieur, Sud-Est de la France) : conséquence taxinomique et phylétiques.- *Annales du Muséum d'Histoire Naturelle de Nice*, t. XXI, p. 179-253.
- BERT D., DELANOY G. & BERSAC S. (2008).- Propositions pour un nouveau découpage biozonal ammonitique, et nouveaux biohorizons pour le Barrémien supérieur.- *Carnets de Géologie / Notebooks on Geology*, Brest, Article 2008/03 (CG2008\_A03), 18 p.
- BERT D., DELANOY G. & CANUT L. (2009).- L'origine des *Imerites* ROUCHADZÉ, 1933 : résultat d'une innovation chez les Gassendiceratinae BERT, DELANOY & BERSAC, 2006 (Ammonoidea, Ancyloceratina).- *Annales de Paléontologie*, Paris, vol. 95, p. 21-35.
- BERT D., BUSNARDO R., DELANOY G. & BERSAC S. (2010).- Problems in the identity of "*Crioceras*" *barremense* KILIAN, 1895 (Ancyloceratida, Late Barremian), and their proposed resolution.- *Carnets de Géologie / Notebooks on Geology*, Brest, Article 2010/01 (CG2010\_A01), 17 p.
- DELANOY G. (1990).- Précisions nouvelles sur "*Emericiceras*" *collignoni* THOMEL (Ammonoidea) et proposition d'un nouveau schéma zonal dans le Barrémien supérieur du Sud-Est de la France.- *Comptes Rendus de l'Académie des Sciences*, Paris, (Série II, Sciences de la Terre et des Planètes), t. 311, p. 259-264.
- DELANOY G. (1995).- Les biozones à Feraudianus, Giraudi et Sarasini du Barrémien supérieur de la région stratotypique d'Angles-Barrême-Castellane (Sud-Est de la France).- *Géologie Alpine*, Grenoble, Mémoire H.S., n° 20 (1994), p. 279-319.
- DELANOY G. (1997).- Biostratigraphie haute résolution du Barrémien supérieur du Sud-Est de la France.- *Comptes Rendus de l'Académie des Sciences*, Paris, (Série II, Sciences de la Terre et des Planètes), t. 325, p. 689-694.
- DELANOY G. (1998).- Biostratigraphie des faunes d'Ammonites à la limite Barrémien-Aptien dans la région d'Angles-Barrême-Castellane. Étude particulière de la Famille des Heteroceratidae (Ancyloceratina, Ammonoidea).- *Annales du Muséum d'Histoire Naturelle de Nice*, t. XII (1997), 270 p.
- DELANOY G. & BERT D. (2006).- Nouvelles unités taxinomiques de la famille des Heteroceratidae SPATH, 1924.- *Annales du Muséum d'Histoire Naturelle de Nice*, t. XXI, p. 255-295.
- DELANOY G. & EBBO L. (2000).- Une nouvelle espèce d'*Heteroceras*: *H. mascarelli* sp. nov. (Ancyloceratina, Ammonoidea) dans le Barrémien supérieur du S-E de la France.- *Annales du Muséum d'Histoire Naturelle de Nice*, t. XV, p. 1-17.
- ERISTAVI M.S. (1955).- Nizhnelovaja fauna Gruzii (Lower Cretaceous fauna of Georgia).- *Institut Geologii i Mineralologii Akademii Nauk G.S.S.R., Monografii*, Tbilisi, 6, 224 p. (en russe).
- GAUTHIER H. avec la collaboration de BUSNARDO R., COMBÉMOREL R., DELANOY G., FISCHER J.-C., GUÉRIN-FRANIATTE F., JOLY B., KENNEDY W.J., SORNAY J. & TINTANT H. (eds.) (2006).- Révision critique de la Paléontologie Française d'Alcide d'ORBIGNY, Vol. IV, Céphalopodes Crétacés, sous la direction de J.C. FISCHER.- Backhuys Publisher, Leiden, 292 p.
- HOEDEMAEKER P.J. & BULOT L. (reporters), AVRAM E., BUSNARDO R., COMPANY M., DELANOY G., KAKABADZE M., KOTETISHVILI E., KRISHNA J., KVANTALIANI I., LATIL J.L., MEMMI L., RAWSON P.F., SANDOVAL J., TAVERA J.M., THIEULOY J.-P., THOMEL G., VASICEK Z. & VERMEULEN J. (1990).- Preliminary ammonite zonation for the lower Cretaceous of Mediterranean region (Lower Cretaceous Cephalopod team, Working Group of IGCP, Project 262).- *Géologie Alpine*, Grenoble, t. 66, p. 123-127.
- KAKABADZE M.V. (1989).- The Barremian biostratigraphical subdivisions of Georgia and comparison with some western mediterranean regions. *In*: WIEDMANN J. (ed.), Cretaceous of the Western Tethys.- Proceedings 3rd International Cretaceous Symposium, Tubingen (1987), E. Schweizerbart'sche Verlagsbuchhandlung, Stuttgart, p. 551-560.
- KAKABADZE M.V. (2004).- Intraspecific and intra-

- generic variabilities and their implication for the systematics of the Cretaceous heteromorph ammonites; a review.- *Scripta Geologica*, Leiden, vol. 128, p. 17-37.
- KILIAN W. (1888).- Sur quelques fossiles du Crétacé inférieur de la Provence.- *Bulletin de la Société Géologique de France*, Paris, (3ème série), t. XVI, p. 663-691, pls. XVII-XXXI.
- KILIAN W. (1889).- Description géologique de la montagne de Lure.- Thèse, Masson édit., Paris, 458 p.
- KILIAN W. (1907-1913).- Das Mesozoicum Unterkreide (Palaeocretacium).- *Lethaea Geognostica*, Stuttgart, Teil III, Bd. 3, p. 234-279.
- KILIAN W. & REBOUL P. (1915).- I. La faune de l'Aptien inférieur des environs de Montélimar (Drôme) (Carrière de l'Homme d'Armes). In: KILIAN W. (ed.), Contribution à l'étude des faunes paléocrétacées du Sud-Est de la France.- *Mémoires pour servir à l'Explication de la Carte Géologique détaillée de la France*, Paris, n° 14, 221 p.
- KOTETISHVILI E.V., KVANTALIANI I.V., KAKABADZE M.V. & TSIREKIDZE L.R. (eds., 2005).- Atlas of Early Cretaceous fauna of Georgia.- *Proceedings of the Georgian Academy of Sciences, A. Janelidze Geological Institute*, Tbilisi, (n. ser.), vol. 120, 788 p.
- MEEK F.B. (1876).- A report on the invertebrate Cretaceous and Tertiary fossils of the Upper Missouri Country.- Report of the United States Geological Survey of the Territories, Washington, vol. IX, 629 p.
- ORBIGNY A. d' (1840-1842).- Paléontologie française. Description des mollusques et rayonnés fossiles - Terrains crétacés. Tome Premier. Céphalopodes.- Masson, Paris, 662 p.
- ORBIGNY A. d' (1851).- Notice sur le genre *Heteroceras*, de la classe des Céphalopodes.- *Journal de Conchyliologie*, Paris, tome second, p. 217-222.
- PAQUIER M.V. (1900).- Recherches géologiques dans le Diois et les Baronnies orientales.- *Travaux du Laboratoire de Géologie de la Faculté des Sciences de l'Université de Grenoble*, t. 5, p. 149-556.
- REBOULET S. (reporter), ATROPS F., BERT D., BULOT L., BUSNARDO R., DELANOY G. & VERMEULEN J. (2007, unpublished).- Zonation Hauterivien – Barrémien.- Compte rendu de la réunion des biostratigraphes français du KILIAN Group (IUGS Lower Cretaceous Ammonites Working Group), Digne-les-Bains (2 Mai 2007), 14 p.
- REBOULET S. & HOEDEMAEKER P.J. (reporters), AGUIRRE URRETA M.B., ALSÉN P., ATROPS F., BARABOSKIN E.Y., COMPANY M., DELANOY G., DUTOUR Y., KLEIN J., LATIL J.L., LUKENEDER A., MITTA V., MOURGUES F.A., PLOCH I., RAISSOSADAT N., ROPOLLO P., SANDOVAL J., TAVERA J.M., VAŠÍČEK Z., VERMEULEN J., ARNAUD H., GRANIER B. & PREMOLI-SILVA I. (2006).- Report on the second international meeting of the IUGS Lower Cretaceous Ammonite Working Group, the "KILIAN Group" (Neuchâtel, Switzerland, 8 September 2005).- *Cretaceous Research*, London, vol. 27, n° 5, p. 712-715.
- REBOULET S. & KLEIN J. (reporters), BARRAGAN R., COMPANY M., GONZALEZ-ARREOLA C., LUKENEDER A., RAISSOSADAT S.N., SANDOVAL J., SZIVES O., TAVERA J.M., VAŠÍČEK Z. & VERMEULEN J. (2009).- Report on the 3rd International Meeting of the IUGS Lower Cretaceous Ammonite Working Group, the "KILIAN Group" (Vienna, Austria, 15th April 2008).- *Cretaceous Research*, London, vol. 30, n° 2, p. 496-502.
- ROUCHADZÉ J. (1933).- Les ammonites aptiennes de la Géorgie occidentale.- *Bulletin de l'Institut géologique de Géorgie*, Tbilisi, vol. 1 (1932), n° 3, p. 165-273.
- SALVADOR A. (ed., 1994).- International stratigraphic guide. A guide to stratigraphic classification, terminology and procedures.- I.U.G.S. & Geological Society of America, 214 p.
- SARKAR S.S. (1955).- Révision des Ammonites déroulées du Crétacé inférieur du SE de la France.- *Mémoires de la Société Géologique de France*, Paris, (N.S.), t. XXXIV, n° 72, 176 p.
- THIERRY J. (1997).- La biostratigraphie, du taxon à la biozone et aux échelles biozonales. In: REY J. (éd.), Stratigraphie, Terminologie Française.- *Bulletin des Centres de Recherches Exploration-Production elf-aquitaine*, Pau, Mémoire 19, p. 63-89.
- VERMEULEN J. (2000).- Nouvelles données sur les répartitions stratigraphiques, les évolutions et les classifications de trois familles d'ammonites du Crétacé inférieur.- *Géologie Alpine*, Grenoble, n° 75, p. 123-132.
- VERMEULEN J. (2006).- Nouvelle classification à fondement phylogénétique des ammonites hétéromorphes du Crétacé inférieur.- *Annales du Muséum d'Histoire Naturelle de Nice*, t. XXI, p. 137-178.
- VERMEULEN J. & LEPINAY P. (2010).- Nouvelles données sur le genre *Imerites* ROUCHADZÉ, 1933 (Ammonidita, Ancyloceratina) ; réponses et précisions apportées au travail de BERT, DELANOY et CANUT, 2009.- *Annales de Paléontologie*, Paris, vol. 96, p. 13-23.
- WRIGHT C.W., CALLOMON J.H. & HOWARTH M.K. (eds., 1996).- Cretaceous Ammonoidea. In: KAESLER R.L. (ed.), Mollusca 4 (revised).- *Treatise on Invertebrate Paleontology*, Geological Society of America, New York; University of Kansas, Lawrence, Part L, 362 p.