

***Heteroporella ? paucicalcare* (CONRAD, 1970), an Urgonian Dasycladalean alga revisited**

Bruno GRANIER ¹

Abstract: When the species *Heteroporella ? paucicalcare* (CONRAD, 1970) was erected it was left in open nomenclature. Later on new combinations were introduced but did not meet a general agreement among paleophycologists. Considering the current acception of the Dasycladalean families, we ascribe it to the Family Polyphysaceae and subsequently to the genus *Clypeina* (MICHELIN, 1845). Its known stratigraphic range is rather brief (Late Hauterivian-Early Barremian), which makes it a good index fossil in Urgonian carbonate platform series. Its geographical distribution is apparently restricted to Western Europe (France, Spain and Switzerland).

Key Words: Calcareous fossil algae; Dasycladales; Polyphysaceae; *Heteroporella*; *Clypeina*; Hauterivian; Lower Barremian.

Citation : GRANIER B. (2013).- *Heteroporella ? paucicalcare* (CONRAD, 1970), an Urgonian Dasycladalean alga revisited.- [Carnets de Géologie \[Notebooks on Geology\]](#), Brest, Letter 2013/01 (CG2013_L01), p. 59-65.

Résumé : Révision de l' *Heteroporella ? paucicalcare* (CONRAD, 1970), une algue dasycladale urgonienne.- À sa création l'espèce *Heteroporella ? paucicalcare* (CONRAD, 1970) a été laissée en nomenclature ouverte. Ultérieurement de nouvelles combinaisons ont été proposées mais aucune n'a rencontrée l'unanimité parmi les paléophycologues. Compte-tenu de l'acception actuelle des familles d'algues dasycladales, nous rapportons ce taxon à la Famille des Polyphysaceae et par contre-coup au genre *Clypeina* (MICHELIN, 1845). Sa répartition stratigraphique connue est plutôt courte (Hauterivien supérieur-Barrémien inférieur), ce qui en fait un bon marqueur fossile dans les séries de plates-formes carbonatées urgoniennes. Sa distribution géographique est apparemment réduite à l'Europe occidentale (Espagne, France et Suisse).

Mots-Clefs : Algues calcaires fossiles ; Dasycladales ; Polyphysaceae ; *Heteroporella* ; *Clypeina* ; Hauterivien ; Barrémien inférieur.

1. Introduction

Further to the 1994 revision of *Heteroporella lepina* PRATURLON, 1967, *i.e.*, the type-species of the genus *Heteroporella* (CROS & LEMOINE ex PRATURLON, 1967), the latter was left with one species only (its type, by definition) out of its original sixteen representatives (GRANIER *et al.*, 1994). Most species were ascribed either to *Chinianella* OTT ex GRANIER & DELOFFRE, 1993, or to *Otternstella* GRANIER in GRANIER *et al.*, 1994; few specific epithets were considered *nomina nuda* whereas five species were left in open nomenclature, awaiting revision. *Heteroporella ? paucicalcare* CONRAD, 1970, is part of this last set and required our attention.

2. Taxonomical background

From its very beginning (CONRAD, 1970), *Heteroporella ? paucicalcare* was in open nomenclature. The question mark following the generic name was fully justified because it was not obvious whether the illustrated specimens were bearing both fertile (large) and sterile (thin)

laterals as stated in the original diagnosis of the genus (PRATURLON, 1967), that is prior to its revision (GRANIER *et al.*, 1994). Actually the pores corresponding to the so-called "sterile hairs, located within and/or between the whorls", which were reported by CONRAD (1970), are merely defined in the interverticillar space by the incompleteness of the calcification on the whorls of fertile laterals.

On the basis of material questionably referred to the species, BUCUR (2000) introduced the new combination, *Similiclypeina paucicalcare* nov. comb. An option that was sound because the genus *Similiclypeina* BUCUR, 1993, was erected to group *Clypeina*-like species the verticils of which are set rather close together ("les espèces clypeiniformes dont les verticilles sont relativement rapprochés", BUCUR, 1994).

But, in turn, CONRAD *et al.* (2009) introduced another new combination, *Piriferella paucicalcare* nov. comb., which was fully backed by BUCUR (*in* CONRAD *et al.*, 2009; BUCUR, 2011). There are some side consequences:

¹ Département des Sciences de la Terre et de l'Univers, UFR Sciences et Techniques, Université de Bretagne Occidentale, 6 Avenue Le Gorgeu, CS 93837, F-29238 Brest Cedex 3 (France)
bgranier@univ-brest.fr

Manuscript online since February 28, 2013

[Editor: Michel MOULLADE; language editor: Donald E. OWEN]

1- for instance, the species *Clypeina somalica* CONRAD *et al.*, 1983, was successively transferred to *Holosporella* PIA, 1930 (GRANIER *et al.*, 1991), then to *Similiclypeina* BUCUR, 1993 (BUCUR, 1993), and finally to *Piriferella* SOKAČ, 1996 (SOKAČ, 1996; CONRAD *et al.*, 2009). CONRAD *et al.* (1983) stated that its thallus bears verticils with noncontiguous branches, even near to the main axis ("*Rameaux (...) jamais contigus même à proximité du siphon*", *ibid.*); accordingly this species should have been excluded from genera *Clypeina* (MICHELIN, 1845) and *Similiclypeina*. The two remaining genera, *Holosporella* PIA, 1930, and *Piriferella* SOKAČ, 1996, have similar characteristics, the only difference between them being the shape of the laterals. Though the species *somalica* better fits the diagnostic parameters of *Holosporella* (GRANIER *et al.*, 1991; GRANIER & DELOFFRE, 1993), CONRAD *et al.* (2009) favoured the *Piriferella* option. In addition, without giving much consideration to the measurements (particularly the spacing of the verticils and the relative size of the fertile pores), these authors also regarded *Piriferella spinosa* SOKAČ, 1996, the type-species, and *Salpingoporella verrucosa* SOKAČ, 1996, as junior synonyms of *Piriferella somalica* (CONRAD *et al.*, 1983), a statement we should not agree with;

2- the genus *Similiclypeina* BUCUR, 1993, is downsized. Actually it might even be considered as monospecific due to the so far unique morphological trait of the proximal part of the laterals in its type-species, *Similiclypeina conradi* BUCUR, 1993;

3- Rather than clearing up the confusion that was there CONRAD *et al.* (2009) added more. Let us remember that *Holosporella* PIA, 1930, *Piriferella* SOKAČ, 1996, and *Salpingoporella* PIA in TRAUTH, 1917, belong to the Family Triploporellaceae (PIA, 1920), *Heteroporella* (CROS & LEMOINE ex PRATURLON, 1967) is a representative of the Family Thyrsoporellaceae (GRANIER & BUCUR in GRANIER *et al.*, 2012), whereas both *Clypeina* (MICHELIN, 1845) and *Similiclypeina* BUCUR, 1993, are ascribed to the Family Polyphysaceae (KÜTZING, 1841).

In conclusion, except for one small detail, it would be safer to revert to BUCUR's (2000) view: *Heteroporella ? paucicalcareia* is a *Clypeina*-like species the verticils of which are set rather close together. Because the species lacks the typical pattern of the laterals in *Similiclypeina conradi* BUCUR, 1993, it should "naturally" be relocated among the representatives of *Clypeina* (MICHELIN, 1845).

3. Description of the new material

Material: The studied material consists of some tens of thin sections, each containing one to several sections (mostly oblique and more or less well preserved) of the alga. Part of the material (Fig. 1), the one from Haute-Savoie (France), is derived from rock samples collected by TRABOLD (1996) during field work to complete a PhD thesis: a first calcareous alga, *Falsolikella danilovae* (RADOIČIĆ ex BARATTOLO, 1978), was already revised based on the same set of thin sections (GRANIER *et al.*, 2000). Additional material (Fig. 2) was collected by the author from a section in Drôme (France), where algal remains transported in debris and grain flows can be dated by the ammonites found in the adjacent marls and limestones (GRANIER *et al.*, nearing completion).

Measurements	CONRAD, 1970	This work (TRABOLD Collection)
L	2.5 mm	> 1.75 mm
D	0.85 - 1.5 mm	0.89 - 1.28 mm
d	0.3 - 0.57 mm	0.32 - 0.47 mm
d/D	30% - 48%	25% - 35%
h1 or h2	0.17 - 0.28 mm (h1)	0.23 - 0.43 mm (h2)
l		0.30 - 0.43 mm
p		0.08 - 0.15 mm
e		0.04 - 0.07 mm
w	9 - 18	8 - 14

Table 1: Biometric data of *Clypeina paucicalcareia* (CONRAD, 1970) *nov. comb.* (L: maximum length; D: external diameter; d: diameter of the stem; h1: interverticillar spacing *sensu stricto*, *i.e.*, from the top surface of one whorl to the bottom surface of the next; h2: interverticillar spacing *sensu lato*, *i.e.*, distance from a reference plane in one whorl to the same plane in the next; l: length of the laterals; p: width of the laterals; e: thickness of the calcareous coating on the laterals; w: number of laterals per verticil).




Figure 1: a-c and e-m: *Clypeina paucicalcareia* (CONRAD, 1970) *nov. comb.*

a. oblique to tangential section. GT 1.065 (Upper Hauterivian, Ha6 HST) Rocher de Cluses, Haute-Savoie, France; **b.** oblique to axial section. GT 1.122 (Upper Hauterivian, Ha7 LST) Rocher de Cluses; **c.** oblique to transverse section. FL 2.09 (Lower Barremian, Ba1 LST) Flaine, Haute-Savoie, France; **e.** axial section. GT 1.066 (Upper Hauterivian, Ha6 HST) Rocher de Cluses; **f.** tangential section. GT 1.047 (Upper Hauterivian, Ha6 TST) Rocher de Cluses; **g.** oblique section through 3 verticils. GT 1.067 (Upper Hauterivian, Ha6 HST) Rocher de Cluses; **h.** oblique section. GT 1.098 (Upper Hauterivian, Ha7 LST) Rocher de Cluses; **i.** subtransverse section. BA 19a05 (Upper Hauterivian, Ha6 LST) Combe de Balme, Haute-Savoie, France; **j.** oblique to transverse section. GT 1.101 (Upper Hauterivian, Ha7 LST) Rocher de Cluses; **k.** oblique section through 3 verticils. GT 1.066 (Upper Hauterivian, Ha6 HST) Rocher de Cluses; **l.** oblique section through 4 verticils. FL 2.09 (Lower Barremian, Ba1 LST) Flaine; **m.** oblique section through (?) 2 verticils. GT 1.122 (Upper Hauterivian, Ha7 LST) Rocher de Cluses.

d: oblique to tangential section of a "look-alike" Triploporellacean alga (*Clypeina* is a Polyphysacean alga). Note that laterals are arranged in quinconces. C5.709 (Upper Hauterivian, Ha7 LST) Plateau d'Andey, Haute-Savoie, France.

All thin sections are stored at the Muséum d'Histoire naturelle, Genève [Scale bar: 250µm]: G. TRABOLD (a-c and e-m) and J. CHAROLLAIS (d) collections [Scale bar: 250µm].




Figure 2: a-e: *Clypeina paucicalcareia* (CONRAD, 1970) *nov. comb.* L'Estellon section, Drôme, France. Ages are directly constrained by ammonites found in strata below and above.

a. tangential section through 4 verticils. EST 26.5 (Nicklesi Zone, Lower Barremian, LST Ba1); **b.** oblique section through 5 verticils. EST 92.3 (Moutonianum Zone, Lower Barremian, LST Ba3); **c.** transverse section of a verticil with 8 primary laterals. EST 49.5 (Pulchella Zone, Lower Barremian, LST Ba2); **d.** oblique section through 3 verticils. EST 92.3 (Moutonianum Zone, Lower Barremian, LST Ba3); **e.** oblique section through 4 verticils. EST 49.7 (Pulchella Zone, Lower Barremian, LST Ba2).

All thin sections from the author's collection [Scale bar: 250μm].

Description: The algal thallus is roughly cylindrical with a cylindrical main axis bearing whorls (euspondyle type) consisting of 8 to 16 (w) laterals. The calcareous skeleton is broken into segments with rarely more than ten verticils. These verticils are close set, but not imbricated. The interverticillar spacing *sensu stricto* (h1), *i.e.*, the distance from the top surface of one whorl to the bottom surface of the next whorl, is always a positive value (see Table 1). The interverticillar spacing *sensu lato* (h2), *i.e.*, the distance from a reference plane in one whorl (insertion points of a verticil, for instance) to the same plane in the next whorl, is more than twice the thickness of a verticil (see Table 1), that is more than twice the diameter of the laterals (p). These laterals are of the first order only, slightly phloiophorous in shape, *i.e.*, increasing in diameter outward, and supposedly

fertile because they form quite wide pores. Within a single verticil, they are first gently inclined (probably upward, in the growth direction) with respect to the algal main axis (up to ? 60°), then they rapidly bend outward and are nearly perpendicular in the medial and distal parts of the whorl, which are almost planar. These laterals are close set in the proximal part of the verticil itself and thus embedded in a joint calcareous coating. But, because their diameter does not increase significantly, they rapidly diverge distally and as a result each lateral gets an individual coating. The star pattern of the verticils in transverse sections (Fig. 2.c) reminds us that it is known in *Clypeina* and in *Actinoporella* (GÜMBEL *in* ALTH, 1881). The lack of corona structure above or below the verticil allows us to ascribe the species *paucicalcareia* to the genus *Clypeina*.

4. Systematics

Phylum Chlorophyta
Class Dasycladophyceae
HOEK *et al.*, 1995
Order Dasycladales PASCHER, 1931
Family Polyphysaceae (KÜTZING, 1841)
Genus *Clypeina* (MICHELIN, 1845)
***Clypeina paucicalcareae* (CONRAD, 1970)**
nov. comb.

(Figs. 1 - 2 - 3)

- 1970** *Heteroporella* ? *paucicalcareae* n. sp.- CONRAD, p. 68-69, Fig. 5 (holotype, duplicated in Fig. 3 herein), Pl. III, figs. 1-4; Pl. IV, figs. 1-3
- 1970** *Heteroporella* ? aff. *paucicalcareae* n. sp.- CONRAD, p. 68-69, Pl. IV, fig. 4
- 1973** *Heteroporella* (?) *paucicalcareae*- JAFFREZO, p. 80, Pl. 3, figs. 9-11 & 16
- 1976** *Heteroporella* ? *paucicalcareae*- PEYBERNÈS, Pl. XXIV, figs. 14-16
- 1976** *Heteroporella* ? *paucicalcareae*- CONRAD & PEYBERNÈS, p. 185-186, Figs. 10.a-b & 13a
- 1976** *Heteroporella* (?) *paucicalcareae*- MASSE, p. 177, Pl. 4, fig. 7
- 1978** *Heteroporella* ? *paucicalcareae*- BASSOULLET *et al.*, p. 133, Pl. 15, figs. 5 (= Fig. 5 in CONRAD, 1970), 6 (= Pl. III, fig. 1 in CONRAD, 1970), 7 (= Fig. 10.a in CONRAD & PEYBERNÈS, 1976) & 8 (= Fig. 10.b in CONRAD & PEYBERNÈS, 1976)
- 1980** *Heteroporella* ? *paucicalcareae*- ARNAUD-VANNEAU, Pl. 111, figs. 3-5
- 1980** *Heteroporella* (?) *paucicalcareae*- JAFFREZO, p. 252-253, Pl. XIX, figs. 1, 2 (= Pl. 3, fig. 9 in JAFFREZO, 1973), 3 & 4 (= Pl. 3, fig. 11 in JAFFREZO, 1973)
- 1989** *Heteroporella* ? *paucicalcareae*- CONRAD & MASSE, p. 281-282, Pl. II, fig. 11
- 1993** *Heteroporella* ? *paucicalcareae*- MASSE, Pl. 2, fig. 1 (= Pl. 4, fig. 7 in MASSE, 1976)
- 1993** *Heteroporella* ? *paucicalcareae*- GRANIER & DELOFFRE, p. 33
- non 1993** *Heteroporella* ? aff. *paucicalcareae*- BODROGI *et al.*, p. 64, Pl. 3, fig. 3
- non 1993** *Heteroporella* (?) *paucicalcareae*- BUCUR *et al.*, Pl. II, fig. 4
- non 1993** *Heteroporella* ? *paucicalcareae*- SOTÁK & MIŠÍK, Pl. 5, figs. 1-2
- 1994** *Heteroporella* ? *paucicalcareae*- GRANIER *et al.*, p. 135
- non 1994** *Heteroporella* ? *paucicalcareae*- BUCUR, p. 152, Pl. VI, fig. 13
- non 2000** *Similiclypeina paucicalcareae* nov. comb.- BUCUR, p. 60, Pl. IV, fig. 7
- non 2000** *Similiclypeina* aff. *paucicalcareae*- BUCUR *et al.*, Pl. VIII, figs. 3-4
- 2007** *Similiclypeina paucicalcareae*- BUCUR *et al.*, Figs. 4.6 & 4.9
- 2007** *Salpingoporella genevensis*- CLAVEL *et al.*, Pl. 6, fig. o
- 2007** *Piriferella paucicalcareae*- CLAVEL *et al.*, Pl. 6, fig. p
- non 2007** *Piriferella paucicalcareae*- CLAVEL *et al.*, Pl. 6, fig. q
- 2009** *Piriferella paucicalcareae* nov. comb.- CONRAD *et al.*, p. 26
- 2011** *Piriferella paucicalcareae*- BUCUR, p. 628, Pl. 1, figs. 6-7; Pl. 2, fig. 13; Pl. 6, figs. 5 & 8

Emended diagnosis: *Clypeina* representative with close set verticils consisting of 8 to 16 slightly phloiophorous laterals. Corresponding pores first gently inclined, then rapidly bending outward and almost horizontal distally. Biometric measurements (Table 1) might help discriminating this species from the other representatives of the genus.



Figure 3: Holotype of *Clypeina paucicalcareae* (CONRAD, 1970) *nov. comb.* Oblique section through 6 verticils. Sample CONRAD 319, slide 6 (Upper Hauterivian, TST Ha5); Rocher-des-Hirondelles, La Rivière, Ain, France; M.A. CONRAD Collection, Muséum d'Histoire naturelle, Genève. Scale bar: 250µm [Some rights reserved].

5. Discussion

Clypeina paucicalcareae (CONRAD, 1970) *nov. comb.* has been reported from strata ranging in age from the Hauterivian to the Bedoulian: (? Upper) Hauterivian (CONRAD & MASSE, 1989), Lower Barremian (CONRAD, 1970; PEYBERNÈS, 1976; CONRAD & PEYBERNÈS, 1976; MASSE, 1976, 1993; ARNAUD-VANNEAU, 1980; BUCUR *et al.*, 2007; CLAVEL *et al.*, 2007), undifferentiated Barremian (JAFFREZO, 1980), undifferentiated Barremian-Bedoulian (JAFFREZO, 1973). The authors reporting the early finds were not able to discriminate the age of the Urganian strata they were dealing with. Today the known range can be reduced to the (?) Late Hauterivian - Early Barremian interval. The first occurrence is not accurate because we lack information on Lower Hauterivian (and even Valanginian) shallow-

water carbonates. The last occurrence is better approximated. According to MASSE (1993) the species is not known above the Pulchella Zone (Early Barremian) although according to CLAVEL (personal communication, Dec. 2012) it is still present in the Moutonianum Zone, *i.e.*, in the last ammonite zone of the Early Barremian.

This Urganian species was found in several localities of W Switzerland (Genève: CONRAD, 1970), S France (Eastern Pyrenees: JAFFREZO, 1973, 1980, PEYBERNÈS, 1976, CONRAD & PEYBERNÈS, 1976; Provence: MASSE, 1976, 1993; Ardèche and Drôme: ARNAUD-VANNEAU, 1980, BUCUR *et al.*, 2007, CLAVEL *et al.*, 2007, BUCUR, 2011), and N Spain (Eastern Pyrenees: CONRAD & PEYBERNÈS, 1976, not illustrated). There it is commonly found associated to the classical "lower" Urganian species: *Falsolikanella danilovae* (RADOIČIĆ *ex* BARATTOLO, 1978), *Pseudoactinoporella fragilis* CONRAD, 1970, *Salpingoporella genevensis* CONRAD *ex* CONRAD *et al.*, 1973, *S. polygonalis* SOKAČ, 1996, *S. muehlbergii* (LORENZ, 1902), *S. melitae* RADOIČIĆ, 1967, ... Records outside that area, *i.e.*, in Hungary (BODROGI *et al.*, 1993), Romania (BUCUR *et al.*, 1993; BUCUR, 1994, 2000), Slovakia (SOTÁK & MIŠÍK, 1993) and Turkey (BUCUR *et al.*, 2000), are not certified. Therefore the MASSE's (1993) hypothesis regarding a possible algal provincialism in Western Europe remains valid.

Acknowledgments

The author thanks F. BARATTOLO and I. BUCUR for their review of the original manuscript, J. CHAROLLAIS for having given him access to his collection and to that of G. TRABOLD, and B. CLAVEL for providing information on the stratigraphic position (including stratigraphic sequence) of the illustrated specimens. Field work (at L'Estellon, Drôme) in 2012 was supported by "Carnets de Géologie" Association.

Bibliographic references

- ARNAUD-VANNEAU A. (1980).- Micropaléontologie, paléocéologie et sédimentologie d'une plateforme carbonatée de la marge passive de la Téthys : l'Urganien du Vercors septentrional et de la Chartreuse (Alpes Occidentales).- *Géologie Alpine*, Grenoble, Mémoire 11, 874 p. (115 Pls.).
- BASSOULLET J.-P., BERNIER P., CONRAD M.A., DELOFFRE R. & JAFFREZO M. (1978).- Les Algues Dasycladales du Jurassique et du Crétacé ; révision critique.- *Géobios*, Villeurbanne, Mémoire spécial, n° 2, 330 p.
- BODROGI I., CONRAD M.A. & LOBITZER H. (1993).- Lower Cretaceous Dasycladales from the Villány zone, Southwest Hungary. Biogeographical significance. In: BARATTOLO F., DE CASTRO P. & PARENTE M. (eds.), Studies on fossil benthic algae.- *Bolletino della Società Paleontologica Italiana*, Modena, Volume Speciale 1, p. 59-68 (3 Pls.).
- BUCUR I.I. (1994).- Algues calcaires de la zone de Reșița-Moldova Nouă (Carpathes Méridionales, Roumanie).- *Revue de Paléobiologie*, Genève, vol. 13, n° 1, p. 147-209 (XXII Pls.).
- BUCUR I.I. (2000).- Lower Cretaceous dasyclad algae from Pădurea Craiului massif (Northern Apuseni Mountains, Romania).- *Acta Palaeontologica Romaniaae*, Cluj Napoca, vol. II (1999), p. 53-72 (VI Pls.).
- BUCUR I.I. (2011).- Early Barremian dasycladalean algae from Serre de Bleyton (Drôme, SE France).- *Annalen des Naturhistorischen Museums in Wien*, Serie A, Band 113, p. 619-653 (7 pls.).
- BUCUR I.I., ARNAUD-VANNEAU A. & ARNAUD H. (2007).- Remarks on *Triploporella praturlonii* BARATTOLO 1982 from the Lower Barremian of Southern Vercors (SE France).- *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*, Stuttgart, Band 244, Heft 2, p. 133-141.
- BUCUR I.I., COCIUBA I. & COCIUBA M. (1993).- Microfacies and microfossils in the Upper Jurassic - Lower Cretaceous limestones in the southern part of the Pădurea Craiului mountains.- *Romanian Journal of Stratigraphy*, Bucuresti, vol. 75, p. 33-40 (IX Pls.).
- BUCUR I.I., KOCH R., KIRMACI Z.M. & TASLI K. (2000).- Les algues Dasycladales du Crétacé inférieur (Calcaire de Berdiga) de Kircaova (région de Kale-Gümüşhane, NE Turquie).- *Revue de Paléobiologie*, Genève, vol. 19, n° 2, p. 435-463.
- CLAVEL B., CHAROLLAIS J., CONRAD M., JAN DU CHÊNE R., BUSNARDO R., GARDIN S., ERBA E., SCHROEDER R., CHERCHI A. DECROUEZ D., GRANIER B., SAUVAGNAT J. & WEIDMANN M. (2007).- Dating and progradation of the Urganian limestone from the Swiss Jura to South-East France.- *Zeitschrift der Deutschen Gesellschaft für Geowissenschaften*, Stuttgart, Band 158, Heft 4, p. 1025-1062.
- CONRAD M.A. (1970).- Barremian and Lower Aptian Dasycladaceae in the area surrounding Geneva (Switzerland).- *Geologica romana*, vol. IX, p. 63-100 (XI Pls.).
- CONRAD M.A. & MASSE J.-P. (1989).- Les algues calcaires des formations carbonatées de l'Hauterivien-Barrémien *pro parte* du Jura vaudois et neuchâtelois (Suisse).- *Mémoires de la Société neuchâteloise des Sciences naturelles*, t. XI, p. 277-290 (Pls. I-II).
- CONRAD M.A. & PEYBERNÈS B. (1976).- Hauterivian-Albian Dasycladaceae from Urganian limestones in the French and Spanish eastern Pyrenees.- *Geologica romana*, vol. XV, p. 175-197.
- CONRAD M.A., PEYBERNÈS B. & MASSE J.-P. (1983).- *Clypeina somalica* n. sp. Dasycladale nouvelle du Crétacé inférieur de la Plaque Africaine (Rép. de Somalie, Italie méridionale).- *Annales de la Société géologique du Nord*, Lille, t. CIII, p. 93-96 (Pl. VI).
- CONRAD M.A., SCHLAGINTWEIT F. & BUCUR I.I.

- (2009).- *Piriferella somalica* (Dasycladalean calcareous green algae) from Tithonian deposits in the Northern Calcareous Alps – synonymy followed by an emendation of *Piriferella* SOKAČ 1998 and *Similiclypeina* BUCUR 1993. In: BASSO D., CARAGNANO A., BROCCHI V. & BENZANI F. (eds.), 6th Regional Symposium of IFAA, 1–5 July 2009, Milan - Italy, Abstract book.- *Annali dell'Università degli Studi di Ferrara*, (Museologia Scientifica e Naturalistica), Volume speciale 2009, p. 26 (abstract).
- GRANIER B., BERTHOU P.-Y. & FOURCADE É. (1991).- The Dasycladalean Algae from the Cretaceous of the New World.- *Transactions of the second geological Conference of the geological Society of Trinidad and Tobago*, Port-of-Spain, April 2nd-8th, 1990, p. 178-183.
- GRANIER B., BUCUR I.I. & TRABOLD G. (2000).- *Falsolikanella danilovae* RADOIČIĆ ex BARATTOLO 1978, n. comb., a Diploporacean alga from the Urgonian facies.- *Acta Palaeontologica Romaniaae*, Cluj Napoca, vol. 2 (1999), p. 177-181 (1 pl.).
- GRANIER B. & DELOFFRE R. (1993).- Inventaire des Algues Dasycladales fossiles. II^o partie- Les Algues Dasycladales du Jurassique et du Crétacé.- *Revue de Paléobiologie*, Genève, vol. 12, n^o 1, p. 19-65.
- GRANIER B., DIAS-BRITO D., BUCUR I.I. & TIBANA P. (2012).- *Brasiliporella*, a new mid-Cretaceous dasycladacean genus: the earliest record of the Tribe Batophoreae.- *Facies*, Erlangen. DOI: 10.1007/s10347-012-0312-6
- GRANIER B., MASSE J.-P. & BERTHOU P.-Y. (1994).- *Heteroporella lepina* PRATURLON, 1967, revisited (followed by taxonomic notes on the so-called "*Heteroporella*" species). In: Proceedings of the International Symposium and Field-Meeting "Alpine Algae '93".- *Beiträge zur Paläontologie*, Wien, n^o 19, p. 129-141.
- JAFFREZO M. (1973).- Les algues calcaires du Jurassique supérieur et du Crétacé inférieur des Corbières. Première partie.- *Revue de Micropaléontologie*, Paris, vol. 16, n^o 2, p. 75-88 (3 Pls.).
- JAFFREZO M. (1980).- Les formations carbonatées des Corbières (France) du Dogger à l'Aptien : micropaléontologie stratigraphique, biozonation, paléoécologie, extension des résultats à la Mésogée.- Thèse Doctorat ès Sciences naturelles, Université Pierre et Marie Curie, Paris, xvii + 615 + 23 p. (XXXIII Pls.).
- MASSE J.-P. (1976).- Les calcaires urgoniens de Provence (Valanginien-Aptien inférieur). Stratigraphie, paléontologie, les paléoenvironnements et leur évolution.- Thèse Doctorat ès Sciences, Université d'Aix-Marseille II, 3 vols., 445 p. (60 Pls.).
- MASSE J.-P. (1993).- Early Cretaceous Dasycladalean biostratigraphy from Provence and adjacent regions (South of France, Switzerland, Spain). A reference for Mesogean correlations. In: BARATTOLO F., DE CASTRO P. & PARENTE M. (eds.), Studies on fossil benthic algae.- *Bolletino della Societa Paleontologica Italiana*, Modena, Volume Speciale 1, p. 311-324 (2 Pls.).
- MICHELIN H. (1840-1847).- Iconographie zoophytologique. Description par localités et terrains des Polypiers fossiles de France et pays environnants.- P. Bertrand, Paris, 348 (Texte) + 79 (Planches) p.
- PEYBERNÈS B. (1976).- Le Jurassique et le Crétacé inférieur des Pyrénées franco-espagnoles entre la Garonne et la Méditerranée.- Thèse Doctorat ès Sciences naturelles, Université Paul Sabatier, Toulouse, 459 p. (XLII Pls.).
- PIA J. (1917).- Familie: Dasycladaceae (ENDL.) CRAM. em. In: TRAUTH F. (ed.), Das Eozänvorkommen bei Radstat im Pongau und seine Beziehungen zu den Gleichalterigen Ablagerungen bei Kirchberg am Wechsel und Wimpasing am Leithagebirge.- *Kaiserliche Akademie der Wissenschaften in Wien, Denkschriften*, (mathematisch-maturwissenschaftliche Klasse), Band 95, p. 209-213 (Pl. I).
- PIA J. (1930).- Upper Triassic fossils from the Burmo-Siamese frontier.- A new Dasycladaceae, *Holosporella siamensis* nov. gen., nov. sp., with a description of the allied genus *Aciculella* PIA.- *Records of the Geological Survey of India*, Calcutta, vol. LXIII, part 1, p. 177-181 (Pl. 4).
- PRATURLON A. (1967).- *Heteroporella lepina*, new dasyclad species from Upper Cenomanian - Lower Turonian of Central Apennines.- *Bollettino della Società Paleontologica Italiana*, Modena, vol. 5 (1966), n^o 2, p. 202-205 (Pls. 51-52).
- SOKAČ B. (1996).- Taxonomic review of some Barremian and Aptian calcareous algae (Dasycladales) from the Dinaric and Adriatic karst regions of Croatia.- *Geologica croatica*, Zagreb, vol. 49, n^o 1, p. 1-79 (22 Pls.).
- SOTÁK J. & MIŠÍK M. (1993).- Jurassic and Lower Cretaceous dasycladalean algae from the Western Carpathians. In: BARATTOLO F., DE CASTRO P. & PARENTE M. (eds.), Studies on fossil benthic algae.- *Bolletino della Societa Paleontologica Italiana*, Modena, Volume Speciale 1, p. 383-404 (12 Pls.).
- TRABOLD G. (1996).- Development of the Urgonian limestones in the Delphino-Helvetian realm (Northern Subalpine chains, Haute-Savoie, France). Sedimentology, sequence stratigraphy and biostratigraphy.- *Publications du Département de Géologie et Paléontologie*, Université de Genève, n^o 20, 187 p.